

Topiramate

Recovery-Focused Treatment for Alcohol Use Disorder

How can topiramate help me cut down or stop drinking?

Topiramate can reduce your urge or desire to drink and can help you reduce or stop drinking. This medication can also decrease your desire to continue to drink if you start drinking again.

How does topiramate work?

- Topiramate acts in your brain to reduce your desire to drink alcohol.
- It can help you drink less and help you choose not to drink.

Will I need other forms of treatment?

A successful alcohol treatment program will include a combination of medicine and social support, like Alcoholics Anonymous (AA) meetings, counseling, individual or group therapy, and sometimes hospital treatment.

What do I need to know before starting topiramate?

- Topiramate is used for seizures and migraines and it has also been shown to help treat alcohol use disorder.
- You should not take topiramate if you are pregnant, planning on getting pregnant or are breastfeeding. Topiramate may make birth control less effective.
- People with kidney stones should not take topiramate.
- Topiramate may interact with seizure medications such as phenytoin, valproic acid and carbamazepine.
- Taking medicine for an alcohol use disorder is not substituting one drug for another.

How should I take topiramate?

- The recommended dose of topiramate for the treatment of alcohol use disorder is 200 milligrams (mg) divided twice daily.
- To minimize side effects you will be started on a low dose that will be increased slowly.
- You should drink plenty of fluids while taking topiramate.



What are some side effects of topiramate?

Common Side Effects	What Can I Do if I Experience This?
Dizziness	Rise slowly to prevent falls
Tingling, prickling or numbness of the skin	Avoid irritants such as hot water, tight fitting shoes or clothing
Mental status changes: <ul style="list-style-type: none">• Slowed thoughts or movements• Difficulty concentrating or problems with memory• Speech difficulties	Contact your provider if you feel there are notable worsening signs or symptoms
Weight loss	<ul style="list-style-type: none">• Try to maintain a healthy balanced diet• Keep a record of any significant changes in your weight and appetite

Less common, but can be severe:

Tell your provider immediately if you have any of the following:

- Mood changes such as depressed mood or irritability.
 - Thoughts of harming yourself or others.
- Changes in vision.
- Blood in urine or painful urination.
- Discuss any symptoms that you think may be due to this medication with your healthcare provider.

All medicines can have side effects. Not everyone has side effects though. They usually get better as your body gets used to the new medicine. Talk with your provider or pharmacist if any of the above side effects trouble you.

How long do I have to take topiramate?

- You and your doctor will decide on your treatment plan. Most people take topiramate for at least 6 months and often longer.
- Do not stop your medication without talking to your healthcare provider first.

Veterans Crisis Line

1-800-273-TALK (8255) or Text - 838255



Your provider and pharmacist can help manage any side effects so you can be successful in your recovery!

Contact info: